

Multiple skew orthogonal polynomials and applications

Log-gases in Caeli Australi, Creswick, 4-15 August, 2025
Shi-Hao Li (School of Mathematics, Sichuan University)



四川大學
SICHUAN UNIVERSITY



Facts

- Integrability in random matrix models has been a long-standing interests for studies in mathematical physics

1. Exact computations for correlation functions/Exact solutions for problems in statistical physics

combinatoric models with integrable kernels

- Forrester, Nagao & Rains 06'
- Nagao & Forrester 02'

random involutions
vicious random walkers

Facts

- Integrability in random matrix models has been a long-standing interests for studies in mathematical physics

1. Exact computations for correlation functions/Exact solutions for problems in statistical physics

combinatoric models with integrable kernels

- Forrester, Nagao & Rains 06'
- Nagao & Forrester 02'

[Chapters 5, 6, 10 in the book
"log-gases and random matrices"]

random involutions
vicious random walkers

Facts

- Integrability in random matrix models has been a long-standing interests for studies in mathematical physics

2. Asymptotic/Limiting behavior of correlation kernels and their applications in Painleve/Toda equations

[Chapters 7, 8, 9 in the book
"Log-gases & random matrices"]

- Selberg integral (Chap. 4)
- symmetric function (Chap. 12)
.....

Motivations

Skew orthogonal polynomials in random matrix theory

- The average characteristic polynomials of Orthogonal ensembles and Symplectic ensembles
- Characterization of some two-component log-gas system with different charges

Motivations

Skew orthogonal polynomials in random matrix theory

- The average characteristic polynomials of Orthogonal ensembles and Symplectic ensembles

– also in vicious random walkers, polynuclear growth model & Pfaffian Schur processes ...

- Characterization of some two-component log-gas system with different charges

– with j.p.d.f. $\sim \prod_{j < k}^{N_1} |\alpha_j - \alpha_k| \prod_{j < k}^{N_2} |\beta_j - \beta_k|^4 \prod_{j=1}^{N_1} \prod_{k=1}^{N_2} |\alpha_j - \beta_k|^2 e^{-V_1(\alpha_j) - V_2(\beta_k)}$

$\alpha_j, \beta_k \in \mathbb{R} \mid S'$ [See also Chap. 15 in the book "log-gases and random matrices"]

Motivations

Skew orthogonal polynomials and integrability

- Skew Christoffel-Darboux kernel

The correlation kernel of orthogonal / symplectic ensemble couldn't only be expressed as SOPs, but as a rank one perturbation of unitary ensemble.

[Adler, Forrester, Nagao & van Moerbeke, 99']

$$\tilde{S}_4(x, y) = * S_2(x, y) \Big|_{N \rightarrow 2N} + \cdot P_{2N}(y) \int_x^{+\infty} * P_{2N-1}(t) dt$$

Motivations

Skew orthogonal polynomials and integrability

- Skew Christoffel-Darboux kernel

[Forrester & -L., 20'] discrete symplectic ensemble (discrete linear lattice & q -lattice)

[-L. Shen, Yu & Forrester, 25'] discrete orthogonal ensemble

Some other discrete ensembles could be found at:

On linear lattice: Borodin-Gorin-Guionnet, Gaussian asymptotics of discrete β -ensemble, Publ. Math. IHES 17'

On q -lattice: Olshanski, Macdonald-level extension of beta ensembles and large N transitions, CMP 22'

Motivations

Skew orthogonal polynomials and integrability

- Skew Christoffel-Darboux kernel

- Wave functions for Pfaff lattice

[Adler, Horozov & van Moerbeke, IMRN, 1999]

[Adler & van Moerbeke, Math. Ann., 2002]

[Adler & van Moerbeke, Duke Math J., 2002]

↓
Connection between Toda & Pfaff lattices

→ introduced time evolutions & found Pfaff lattice from SOPS
→ Existence & uniqueness of Pfaffian τ -functions,
Fay identity, and Virasoro constraints

Motivations

Multiple orthogonal polynomials in random matrix theory

- Hermitian matrix ensemble with external source

[Brezin & Hikami, Zinn-Justin, Bleher & Kuijlaars, Desrosiers & Forrester, ...]

- Non-intersecting Brownian motion with different starting point and ending point

Aptekarev, Bleher & Kuijlaars, Daems & Kuijlaars $\rightarrow$ MOPs of mixed type

Motivations

Multiple orthogonal polynomials in classical integrable systems

- Multi-component Toda system

Adler - van Moerbeke - Vanhaecke (CMP 09').

use Cauchy transform to give a whole multi-component Toda hierarchy,
and characterize the gap probability by using the differential equation
w.r.t the end points of intervals ...

Motivations

Multiple orthogonal polynomials in classical integrable systems

- Multi-component Toda system

Álvarez-Fernández, Prieto & Mañas (Adv. Math. 10').

Group decomposition of moment matrix to give dressing operators,

Lax pairs of multi-component Toda system.

References (about random matrices and integrable systems)

Random matrix models	Orthogonality	Integrable lattice hierarchy	References
Hermitian (unitarily invariant) ensemble	Standard orthogonality	1d-Toda lattice/KdV hierarchy	[Gerasimov et al 91'] [Adler&van Moerbeke 95']
Orthogonal/symplectic invariant ensemble	Skew OPs	Pfaff lattice/DKP hierarchy	[Adler, Horozov&van Moerbeke 99'] [Adler&van Moerbeke 02']
Multi-matrix Ensemble	String OPs	2d-Toda hierarchy	[Adler&van Moerbeke 01']
Cauchy two-matrix model	Cauchy bi-orthogonal polynomials	CKP/C-Toda hierarchy	[SHL&Li 19']
Bures ensemble	Partial skew orthogonal polynomials	BKP/B-Toda hierarchy	[Hu&SHL 17'] [Chang, He, Hu&SHL 18']

Questions

- Whether do we have a concept of “Multiple skew-orthogonal polynomials” from random matrix theory or non-intersecting Brownian motions?

Questions

- Whether do we have a concept of “Multiple skew-orthogonal polynomials” from random matrix theory or non-intersecting Brownian motions?
- Is there any connection between multiple skew-orthogonal polynomials and classical integrable systems?

Questions

- Whether do we have a concept of “Multiple skew-orthogonal polynomials” from random matrix theory or non-intersecting Brownian motions?
- Is there any connection between multiple skew-orthogonal polynomials and classical integrable systems?

Reference:

SHL, B. Shen, J. Xiang and G. Yu, Multiple Skew-Orthogonal Polynomials and 2-component Pfaff lattice hierarchy. Ann. Henri Poincare, 25 (2024) 3333-3370.

Skew-orthogonal polynomials

- Definition: Let $S(x, y)$ be a skew symmetric kernel, i.e. $S(x, y) = -S(y, x)$, then one can define a skew inner product $\langle f(x), g(y) \rangle = \int_{\mathbb{R}^2} f(x) S(x, y) g(y) \omega(x) \omega(y) dx dy$. A family of poly. $\{P_n(x)\}$ is called SOPs if

$$\langle P_{2n}(x), P_{2m}(y) \rangle = \langle P_{2n+1}(x), P_{2m+1}(y) \rangle = 0, \quad \langle P_{2n}(x), P_{2m+1}(y) \rangle = h_n \delta_{n,m}, \quad h_n > 0.$$

— OE: $S(x, y) = \operatorname{sgn}(y-x)$, SE: $S(x, y) = \delta'(y-x)$, Bures: $S(x, y) = \frac{y-x}{x+y}$

— valid for discrete measure as well

Skew-orthogonal polynomials

- Definition: $\langle p_{2n}(x), y^i \rangle = 0, \quad i=0, 1, 2, \dots, 2n$
 $\langle p_{2n+1}(x), y^i \rangle = 0, \quad i=0, 1, 2, \dots, 2n+1$

$$p_{2n}(x) = x^{2n} + a_{2n,2n-1}x^{2n-1} + \dots + a_{2n,0} \quad p_{2n+1}(x) = x^{2n+1} + a_{2n+1,2n}x^{2n} + \dots + a_{2n+1,0}$$

- Linear algebra technique and Pfaffian expressions:

$$\begin{pmatrix} m_{0,0} & \dots & m_{0,2n} \\ m_{1,0} & \dots & m_{1,2n} \\ \vdots & & \vdots \\ m_{2n-1,0} & \dots & m_{2n-1,2n} \end{pmatrix} \begin{pmatrix} a_{2n,0} \\ a_{2n,1} \\ \vdots \\ a_{2n,2n-1} \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix} \rightarrow p_{2n}(x) = \frac{1}{T_{2n}} \begin{vmatrix} m_{0,0} & \dots & m_{0,2n-1} & m_{0,2n} \\ \vdots & & \vdots & \vdots \\ m_{2n-1,0} & \dots & m_{2n-1,2n-1} & m_{2n-1,2n} \\ 1 & \dots & x^{2n-1} & x^{2n} \end{vmatrix}$$

$\hookrightarrow m_{i,j} = \langle x^i, y^j \rangle$

Skew-orthogonal polynomials

- Definition: $\langle p_{2n}(x), y^i \rangle = 0, \quad i = 0, 1, 2, \dots, 2n$
 $\langle p_{2n+1}(x), y^i \rangle = 0, \quad i = 0, 1, 2, \dots, 2n+1$

$$p_{2n}(x) = x^{2n} + a_{2n,2n-1}x^{2n-1} + \dots + a_{2n,0} \quad p_{2n+1}(x) = x^{2n+1} + a_{2n+1,2n}x^{2n} + \dots + a_{2n+1,0}$$

- Linear algebra technique and Pfaffian expressions:

$$p_{2n}(x) = \frac{1}{T_{2n}} \text{pf} \begin{pmatrix} m_{0,0} & \dots & m_{0,2n} & -1 \\ \vdots & & \vdots & \vdots \\ m_{2n,0} & \dots & m_{2n,2n} & -x^{2n} \\ 1 & \dots & x^{2n} & 0 \end{pmatrix} := \frac{1}{T_{2n}} \text{pf}(0, \dots, 2n, x)$$

$$\text{pf}(i, j) = m_{ij}, \quad \text{pf}(i, x) = x^i$$

Skew-orthogonal polynomials

- Definition: $\langle P_{2n}(x), y^i \rangle = 0, \quad i = 0, 1, 2, \dots, 2n$
 $\langle P_{2n+1}(x), y^i \rangle = 0, \quad i = 0, 1, 2, \dots, 2n+1$

$$P_{2n}(x) = x^{2n} + a_{2n,2n-1}x^{2n-1} + \dots + a_{2n,0} \quad P_{2n+1}(x) = x^{2n+1} + a_{2n+1,2n}x^{2n} + \dots + a_{2n+1,0}$$

- Linear algebra technique and Pfaffian expressions:

$$\tau_{2n} = \text{pf}(0, 1, \dots, 2n-1) = \frac{1}{(2n)!} \int_{\mathbb{R}^{2n}} \text{pf}(S(x_i, x_j))_{i,j=1}^{2n} \Delta_{2n}(x) \prod_{i=1}^n \omega(x_i) dx_i$$

(de Bruijn's formula, 1955)

Ref.: [Chang, He, Hu & SHL, (Partial)-skew-orthogonal polynomials and related integrable lattices with Pfaffian τ -functions, CMP, 18].

Multiple skew-orthogonal polynomials

- Definition:

Definition 2.3. Given a multi-index $\vec{v} = (v_1, \dots, v_p)$ such that $|\vec{v}| = v_1 + \dots + v_p$ is odd, if there are p different weights $(\omega_1, \dots, \omega_p)$ supported on γ and $\mathbb{S}(x, y)$ is a skew symmetric function from $\gamma \times \gamma$ to $\mathbb{R}$ so that all moments are finite, then for a fixed integer $b \in \{1, 2, \dots, p\}$, there exist multiple skew-orthogonal polynomials $R_1(x), \dots, R_p(x)$ and $\tilde{R}_b(x)$, such that

$$\begin{aligned} \int_{\gamma \times \gamma} \left(\sum_{i=1}^p R_i(x) \omega_i(x) \right) \mathbb{S}(x, y) y^j \omega_k(y) dx dy &= 0, \quad j = 0, \dots, v_k - 1, \quad k = 1, \dots, p, \\ \int_{\gamma \times \gamma} \left(\sum_{i=1}^p R_i(x) \omega_i(x) \right) \mathbb{S}(x, y) \left(\sum_{\substack{i=1 \\ i \neq b}}^p R_i(y) \omega_i(y) + \tilde{R}_b(y) \omega_b(y) \right) dx dy &= 1, \end{aligned} \tag{2.15}$$

where $\deg R_i(x) \leq v_i - 1$ ($i = 1, \dots, p$), and $\deg \tilde{R}_b(x) \leq v_b$. Here, we assume that coefficients in the highest-order terms of R_b and $\tilde{R}_b$ are the same.

Multiple skew-orthogonal polynomials

- Definition:

Definition 2.3. Given a multi-index $\vec{v} = (v_1, \dots, v_p)$ such that $|\vec{v}| = v_1 + \dots + v_p$ is odd, if there are p different weights $(\omega_1, \dots, \omega_p)$ supported on γ and $\mathbb{S}(x, y)$ is a skew symmetric function from $\gamma \times \gamma$ to $\mathbb{R}$ so that all moments are finite, then for a fixed integer $b \in \{1, 2, \dots, p\}$, there exist multiple skew-orthogonal polynomials $R_1(x), \dots, R_p(x)$ and $\tilde{R}_b(x)$, such that

$$\int_{\gamma \times \gamma} \left(\sum_{i=1}^p \underline{R_i(x) \omega_i(x)} \right) \mathbb{S}(x, y) y^j \omega_k(y) dx dy = 0, \quad j = 0, \dots, v_k - 1, \quad k = 1, \dots, p,$$

$$\int_{\gamma \times \gamma} \left(\sum_{i=1}^p \cancel{R_i(x) \omega_i(x)} \right) \mathbb{S}(x, y) \left(\sum_{\substack{i=1 \\ i \neq b}}^p R_i(y) \omega_i(y) + \tilde{R}_b(y) \omega_b(y) \right) dx dy = 1,$$

$R_{(v_1, \dots, v_p)}^{(i)}$ i -th component of polynomials $\tilde{R}_{(v_1, \dots, v_p)}$

(2.15)

where $\deg R_i(x) \leq v_i - 1$ ($i = 1, \dots, p$), and $\deg \tilde{R}_b(x) \leq v_b$. Here, we assume that coefficients in the highest-order terms of R_b and $\tilde{R}_b$ are the same.

Multiple skew-orthogonal polynomials

- Pfaffian expressions:

① A determinantal expression for block matrices

② The use of determinant identity to Pfaffian expressions

$$\begin{aligned} \text{e.g.: } R_{(n_1, n_2)}(x) &= R_{(n_1, n_2)}^{(1)}(x) \omega_1(x) + R_{(n_1, n_2)}^{(2)}(x) \omega_2(x) \\ &= \text{Pf} \begin{bmatrix} A_{n_1, n_1}^{(1,1)} & A_{n_1, n_2}^{(1,2)} & -\psi_1^T(x) \omega_1(x) \\ A_{n_2, n_1}^{(2,1)} & A_{n_2, n_2}^{(2,2)} & -\psi_2^T(x) \omega_2(x) \\ \psi_1(x) \omega_1(x) & \psi_2(x) \omega_2(x) & 0 \end{bmatrix} \end{aligned}$$

Application 1: Non-intersecting Brownian motions

- Karlin-McGregor formula vs LGV (Lindstorm-Gessel-Viennot) lemma

Karlin-McGregor formula: Consider n independent copies with given starting points $a_1 < a_2 < \dots < a_n$ at time $t=0$. Then the probability density of the event of finding the paths at points $x_1, \dots, x_n$ at time $t > 0$ without any two of them having intersected in the time interval $[0, t]$ is proportional to $\det [p(a_i, x_j; t)]_{i,j=1}^n$, where $p(x, y; t)$ is the transition probability density of the diffusion process.

Application 1: Non-intersecting Brownian motions


- Karlin-McGregor formula vs LGV (Lindstorm-Gessel-Viennot) lemma

LGV lemma: Let $\vec{u} = (u^{(1)}, \dots, u^{(r)})$ and $\vec{v} = (v^{(1)}, \dots, v^{(r)})$ be two ordered r -tuples of vertices in an acyclic directed graph D . If $\vec{u}$ and $\vec{v}$ are D -compatible, then

$$\text{GF}[\mathcal{P}_0(\vec{u}; \vec{v})] = \det [h(u^{(i)}, v^{(j)})]_{i,j=1}^r.$$

Application 1: Non-intersecting Brownian motions

- Karlin-McGregor formula vs LGV (Lindstorm-Gessel-Viennot) lemma



$$GF[u^{(1)} \rightarrow v^{(1)}, u^{(2)} \rightarrow v^{(2)}]$$


$$= \omega_1 \omega_6 + \omega_1 \omega_4 \omega_5 + \omega_2 \omega_3 \omega_6$$

$$\det \begin{bmatrix} h(u^{(1)}, v^{(1)}) & h(u^{(1)}, v^{(2)}) \\ h(u^{(2)}, v^{(1)}) & h(u^{(2)}, v^{(2)}) \end{bmatrix}$$

$$= \det \begin{bmatrix} \omega_1 + \omega_2 \omega_3 & \omega_2 \omega_5 \\ \omega_3 \omega_4 & \omega_4 \omega_5 + \omega_6 \end{bmatrix}$$

Application 1: Non-intersecting Brownian motions

- Karlin-McGregor formula vs LGV (Lindstorm-Gessel-Viennot) lemma
- Applications into non-intersecting Brownian motions



$$\mathbb{P} \left[x^{(1)}, \dots, x^{(n)} \mid \begin{array}{l} \text{starts at } z^{(1)}, \dots \\ z^{(n)} \text{ at } t=0; \\ \text{ends at } y^{(1)}, \dots \\ y^{(n)} \text{ at } t=1 \end{array} \right]$$

$$\propto \det [p(\vec{z}, \vec{x}, t_0)] \times$$

$$\det [p(\vec{x}, \vec{y}, 1-t_0)]$$

$$\hookrightarrow p(a, b, t) = \frac{1}{\sqrt{\pi t}} e^{-\frac{(b-a)^2}{2t}}$$

Application 1: Non-intersecting Brownian motions

- Karlin-McGregor formula vs LGV (Lindstorm-Gessel-Viennot) lemma
- Applications into non-intersecting Brownian motions
- Confluent cases and multiple orthogonal polynomials

$$\lim_{\substack{z^{(1)}, \dots, z^{(K)} \rightarrow z_1 \\ z^{(K+1)}, \dots, z^{(n)} \rightarrow z_2}} \det [\mathcal{P}(\vec{z}, \vec{x}, t)] \sim \det \left[\begin{array}{c} x_i^{j-1} \mathcal{P}(z_1, x_i, t) \\ x_i^{l-1} \mathcal{P}(z_2, x_i, t) \end{array} \right]_{\substack{j=1, \dots, K \\ l=1, \dots, n-K}}$$

j.p.d.f. $\propto \Delta_n(x) \det \left[\begin{array}{c} x_i^{j-1} \mathcal{P}(z_1, x_i, t) \\ x_i^{l-1} \mathcal{P}(z_2, x_i, t) \end{array} \right]$ starting from z_1, z_2 and ending at 0

Multiple orthogonal polynomials of type I / II

Application 1: Non-intersecting Brownian motions

- Karlin-McGregor formula vs LGV (Lindstorm-Gessel-Viennot) lemma
- Applications into non-intersecting Brownian motions
- Confluent cases and multiple orthogonal polynomials

$$\lim_{\substack{z^{(1)}, \dots, z^{(K)} \rightarrow z_1 \\ z^{(K+1)}, \dots, z^{(n)} \rightarrow z_2}} \det [\mathcal{P}(\bar{z}, \bar{x}, t)] \sim \det \left[\begin{array}{c} x_i^{j-1} p(z_1, x_i, t) \\ x_i^{l-1} p(z_2, x_i, t) \end{array} \right]_{\substack{j=1, \dots, K \\ l=1, \dots, n-K}}$$

$$\text{j.p.d.f.} \propto \det \left[\begin{array}{c} x_i^{j-1} p(z_1, x_i, t) \\ \vdots \\ x_i^{j_l-1} p(z_l, x_i, t) \end{array} \right] \times \det \left[\begin{array}{c} x_i^{q_1-1} p(x_i, y_1, 1-t) \\ \vdots \\ x_i^{q_m-1} p(x_i, y_m, 1-t) \end{array} \right] \begin{array}{l} \text{starting from } z_1, \dots, z_l \\ \text{ending at } y_1, \dots, y_m \end{array}$$

Multiple orthogonal polynomials of mixed type [Daems & Kuijlaars, 04']


Application 1: Non-intersecting Brownian motions

- Stembridge's formula


Let $\vec{u} = (u^{(1)}, \dots, u^{(2n)})$ be $2n$ -tuple of vertices in an acyclic directed graph $\mathcal{D}$. If $I \subset V$ is a totally ordered subset of vertices s.t. $\vec{u}$ is $\mathcal{D}$ -compatible with I , then $GF[\mathcal{D}_0(\vec{u}; I)] = \text{Pf}[Q_I(u^{(i)}, u^{(j)})]_{i,j=1}^{2n}$,
where $Q_I(u^{(i)}, u^{(j)}) = \sum_{x,y \in I} h(u^{(i)}, x) h(u^{(j)}, y) \text{sgn}(y-x)$.

Application 1: Non-intersecting Brownian motions

- Stembridge's formula



$$\vec{u} \rightarrow \vec{v} \oplus \vec{I}$$




Ref.:

Ciucu and Krattenthaler, The interaction of a gap with a free boundary in a two dimensional dimer system, Commun. Math. Phys., 2011.

Application 1: Non-intersecting Brownian motions

- Stembridge's formula
- Applications into NIBM (distribution of ending points)



$$\mathbb{P} \left[x_t^{(1)}, \dots, x_t^{(n)} \mid \text{starts at } x_0^{(1)}, \dots, x_0^{(n)} \right]$$


$$\propto \det \left[p(\vec{x}_0, \vec{x}_t; t) \right]$$

Ref.:

Katori and Tanemura, Scaling limits of vicious walks and two-matrix models, Phys. Rev. E, 2002.

Application 1: Non-intersecting Brownian motions

- Stembridge's formula
- Applications into NIBM (distribution of ending points)



Ref.:

Katori and Tanemura, Scaling limits of vicious walks and two-matrix models, Phys. Rev. E, 2002.

A discrete version was given by Forrester & Nagao, 2002.

$$\mathbb{P} \left[x_t^{(1)}, \dots, x_t^{(n)} \mid \text{starts at } x_0^{(1)}, \dots, x_0^{(n)} \right] \\ \propto \det \left[p(\vec{x}_0, \vec{x}_t; t) \right]$$

Application 1: Non-intersecting Brownian motions

- A generalization of Stembridge's formula and multiple skew-orthogonal polynomials

Let $\vec{u} = (u^{(1)}, \dots, u^{(r)})$ and $\vec{v} = (v^{(1)}, \dots, v^{(s)})$ be sequences of vertices in an acyclic direct graph $\mathcal{D}$. Assume that I & J are totally ordered subsets of V , s.t. $J \oplus \vec{u}$ and $\vec{v} \oplus I$ are $\mathcal{D}$ -compatible and separated.

If $r, s \in 2\mathbb{N}$, then $GF[\mathcal{D}_o(J \oplus \vec{u}; \vec{v} \oplus I)] = \text{pf} \begin{bmatrix} Q & H \\ -H^T & \tilde{Q} \end{bmatrix}.$


$$Q = [Q_I(u^{(i)}, u^{(j)})], \quad \tilde{Q} = [Q_J(v^{(i)}, v^{(j)})], \quad H = [h(u^{(i)}, v^{(s+1-j)})]$$

Ref.:

SHL and Z. Yao, Non-intersecting path explanation for block Pfaffians and applications into skew-orthogonal polynomials, Adv. Appl. Math., 2025.

Application 1: Non-intersecting Brownian motions

- A generalization of Stembridge's formula and multiple skew-orthogonal polynomials



$$\mathbb{P} \left(\begin{array}{c} \alpha^{(1)}, \dots, \alpha^{(k)} \\ \alpha^{(k+1)}, \dots, \alpha^{(n)} \end{array} \middle| \begin{array}{l} \text{first } k \text{ points starts at } x^{(1)}, \dots, x^{(k)} \\ \text{the rest } n-k \text{ points ends at } x^{(k+1)}, \dots, x^{(n)} \end{array} \right)$$

$$\sim \det \left[p(\vec{\alpha}, \vec{x}; t) \right]$$

Ref.:

SHL and Z. Yao, Non-intersecting path explanation for block Pfaffians and applications into skew-orthogonal polynomials, Adv. Appl. Math., 2025.

Application 1: Non-intersecting Brownian motions

- A generalization of Stembridge's formula and multiple skew-orthogonal polynomials

$$\lim_{\substack{x^{(1)}, \dots, x^{(k)} \rightarrow x_1 \\ x^{(k+1)}, \dots, x^{(n)} \rightarrow x_2}} \det \left[p(\vec{\alpha}, \vec{x}; t) \right]$$

$$\left\{ \begin{array}{l} \left(\begin{array}{c} \alpha^{(1)}, \dots, \alpha^{(k)} \\ \alpha^{(k+1)}, \dots, \alpha^{(n)} \end{array} \middle| \begin{array}{l} \text{first } k \text{ points starts at } x^{(1)}, \dots, x^{(k)} \\ \text{the rest } n-k \text{ points ends at } x^{(k+1)}, \dots, x^{(n)} \end{array} \right) \\ \sim \det \left[p(\vec{\alpha}, \vec{x}; t) \right] \end{array} \right. \leftarrow$$

$$\det \left[\begin{array}{c} \alpha_j^{\ell-1} p(\alpha_j, x_1, t) \\ \alpha_j^{m-1} p(\alpha_j, x_2, t) \end{array} \right]_{\substack{\ell=1, \dots, k \\ m=1, \dots, n-k}}$$

Ref.:

SHL and Z. Yao, Non-intersecting path explanation for block Pfaffians and applications into skew-orthogonal polynomials, Adv. Appl. Math., 2025.

Application 2: 2-component Pfaff lattice

- Time evolutions: Recall: $R_{(n_1, n_2)}(x) = R_{(n_1, n_2)}^{(1)}(x)w_1(x) + R_{(n_1, n_2)}^{(2)}(x)w_2(x)$

$$\omega_1(x; t) = \omega_1(x) \exp\left(\sum_{i=1}^{+\infty} t_i x^i\right) \quad \omega_2(x; t) = \omega_2(x) \exp\left(\sum_{i=1}^{+\infty} s_i x^i\right)$$

$$\begin{cases} \partial_{t_1} (\star R_{(n_1, n_2)}^{(1)}(x; t, s)) = \star' \tilde{R}_{(n_1, n_2)}^{(1)}(x; t, s) \\ \partial_{s_1} (\star R_{(n_1, n_2)}^{(2)}(x; t, s)) = \star' \tilde{R}_{(n_1, n_2)}^{(2)}(x; t, s) \end{cases}$$

Ref.:

1. P. van Moerbeke. Nonintersecting Brownian motions, integrable systems and orthogonal polynomials, MSRI Publ., 2007.
2. M. Adler, P. van Moerbeke, and P. Vanhaecke. Moment Matrices and Multi-Component KP, with applications in random matrix theory, Comm. Math. Phys., 2009.

Application 2: 2-component Pfaff lattice

- Time evolutions: Recall: $R_{(n_1, n_2)}(x) = R_{(n_1, n_2)}^{(1)}(x)w_1(x) + R_{(n_1, n_2)}^{(2)}(x)w_2(x)$

$$\omega_1(x; t) = \omega_1(x) \exp\left(\sum_{i=1}^{+\infty} t_i x^i\right) \quad \omega_2(x; t) = \omega_2(x) \exp\left(\sum_{i=1}^{+\infty} s_i x^i\right)$$

$$\begin{cases} \partial_{t_1} (* R_{(n_1, n_2)}^{(1)}(x; t, s)) = * \tilde{R}_{(n_1, n_2)}^{(1)}(x; t, s) \\ \partial_{s_1} (* R_{(n_1, n_2)}^{(2)}(x; t, s)) = * \tilde{R}_{(n_1, n_2)}^{(2)}(x; t, s) \end{cases}$$

Time evolutions are also imposed on
the partition functions / normalizations / τ -functions

Ref.:

1. P. van Moerbeke. Nonintersecting Brownian motions, integrable systems and orthogonal polynomials, MSRI Publ., 2007.
2. M. Adler, P. van Moerbeke, and P. Vanhaecke. Moment Matrices and Multi-Component KP, with applications in random matrix theory, Comm. Math. Phys., 2009.

Application 2: 2-component Pfaff lattice

- Time evolutions:

Evolutions on wave functions and τ -functions

- Discrete spectral transformations:

$$\begin{aligned} \tau_{(v_1, v_2-1)} d_{(v_1+1, v_2+1)}^{(2)} R_{(v_1+1, v_2+1)}^{(2)}(x) &= \tau_{(v_1+1, v_2)} d_{(v_1, v_2)}^{(2)} \tilde{R}_{(v_1, v_2)}^{(2)}(x) \\ &\quad - \partial_{s_1} \tau_{(v_1+1, v_2)} d_{(v_1, v_2)}^{(2)} R_{(v_1, v_2)}^{(2)}(x) + \tau_{(v_1, v_2+1)} d_{(v_1+1, v_2-1)}^{(2)} R_{(v_1+1, v_2-1)}^{(2)}(x) \end{aligned}$$

Relations between MSOPs with adjacent indexes.

—More examples about discrete spectral transformations of skew-orthogonal polynomials could be found at:

SHL, G. Yu, Christoffel transformations for (partial-)skew-orthogonal polynomials and applications, Adv. Math., 2024.

Application 2: 2-component Pfaff lattice

- Time evolutions:

Evolution on wave functions and τ -functions

Comparison of coefficients of polynomials

Compatible condition of Lax pairs

$$D_{t_1} T_{(n_1, n_2-1)} \cdot T_{(n_1, n_2+1)} = D_{s_1} T_{(n_1+1, n_2)} \cdot T_{(n_1-1, n_2)}$$

$$D_{s_1} D_{t_1} T_{(n_1-1, n_2)} \cdot T_{(n_1-1, n_2)} = 2 \left(T_{(n_1, n_2-1)} T_{(n_1-2, n_2+1)} - T_{(n_1, n_2+1)} T_{(n_1-2, n_2-1)} \right)$$

...

- Discrete spectral transformations:

$$\begin{aligned} \tau_{(v_1, v_2-1)} d_{(v_1+1, v_2+1)}^{(2)} R_{(v_1+1, v_2+1)}^{(2)}(x) &= \tau_{(v_1+1, v_2)} d_{(v_1, v_2)}^{(2)} \tilde{R}_{(v_1, v_2)}^{(2)}(x) \\ &- \partial_{s_1} \tau_{(v_1+1, v_2)} d_{(v_1, v_2)}^{(2)} R_{(v_1, v_2)}^{(2)}(x) + \tau_{(v_1, v_2+1)} d_{(v_1+1, v_2-1)}^{(2)} R_{(v_1+1, v_2-1)}^{(2)}(x) \end{aligned}$$

Relations between MSOPs with adjacent indexes.

—More examples about discrete spectral transformations of skew-orthogonal polynomials could be found at:

SHL, G. Yu, Christoffel transformations for (partial-)skew-orthogonal polynomials and applications, Adv. Math., 2024.

Application 2: 2-component Pfaff lattice

- Cauchy/Stieltjes transform:

$$C_{\psi} g(z) = \int_{\gamma \times \gamma} \frac{\psi(z)}{x-z} S(x,y) g(y) dx dy \Rightarrow$$

$$\begin{aligned} \mathcal{C}_{\omega_1} \left(d_{(v_1, v_2)}^{(i)} R_{(v_1, v_2)}^{(i)} \right) &= (-1)^{v_1} z^{-(v_1+1)} \tau_{(v_1+1, v_2)}(\mathbf{t} + [z^{-1}], \mathbf{s}), \\ \mathcal{C}_{\omega_2} \left(d_{(v_1, v_2)}^{(i)} R_{(v_1, v_2)}^{(i)} \right) &= z^{-(v_2+1)} \tau_{(v_1, v_2+1)}(\mathbf{t}, \mathbf{s} + [z^{-1}]). \end{aligned}$$

- 2-component Pfaff lattice hierarchy:

$$\begin{aligned} & (-1)^{u_1+v_1} \int_{C_{\infty}} e^{\mathfrak{g}(t-t', z)} z^{v_1-u_1-2} \tau_{(v_1-1, v_2)}(t - [z^{-1}], s) \tau_{(u_1+1, u_2)}(t' + [z^{-1}], s') dz \\ & + (-1)^{u_1+v_1} \int_{C_{\infty}} e^{\mathfrak{g}(t-t', z)} z^{u_1-v_1-2} \tau_{(v_1+1, v_2)}(t + [z^{-1}], s) \tau_{(u_1-1, u_2)}(t' - [z^{-1}], s') dz \\ & = \int_{C_{\infty}} e^{\mathfrak{g}(s-s', z)} z^{v_2-u_2-2} \tau_{(v_1, v_2+1)}(t, s - [z^{-1}]) \tau_{(u_1, u_2+1)}(t', s' + [z^{-1}]) dz \\ & + \int_{C_{\infty}} e^{\mathfrak{g}(s-s', z)} z^{u_2-v_2-2} \tau_{(v_1, v_2+1)}(t, s + [z^{-1}]) \tau_{(u_1, u_2+1)}(t', s' - [z^{-1}]) dz \end{aligned}$$

Remarks on Pfaff lattice hierarchy

- The term “Pfaff lattice hierarchy” was first proposed by Adler and van Moerbeke [M. Adler, E. Horozov and P. van Moerbeke, IMRN, 1999]. In fact, this equation is equivalent to the KP equation of D-type according to the classification by Kyoto school.
- The 2-component Pfaff lattice hierarchy was studied by Takasaki [K. Takasaki, Auxiliary linear problem, Difference Fay identities and dispersionless limit of the Pfaff-Toda lattice hierarchy, SIGMA, 2009]. The infinite-dimensional Lie algebraic realization of the general multi-component case is still unknown.
- An elliptic skew orthogonal polynomials and related Pfaff lattice (Landau-Lifshitz equation) was considered recently [W. Fu and SHL, Skew-orthogonal polynomials and Pfaff lattice hierarchy associated with an elliptic curve, IMRN, 2024].



Thanks for your attention!